


Ordering Information
 Web www.encorbio.com
 Email admin@encorbio.com
 Phone 352-372-7022
 Fax 352-372-7066

HGNC Name: CALR
UniProt: P27797
RRID: AB_2572240
Immunogen: Synthetic peptides VESGSLEDDWDFLPPKKI corresponding to amino acids 191-208 of human calreticulin, including the LC3 interacting region or LIR.
Format: Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM NaN₃
Storage: Store at 4°C for short term, for longer term at -20°C
Recommended dilutions:
 WB: 1:1,000-1:2,000 IF/ICC 1:1,000 IHC: 1:500.


References:

- Ostwald TJ, MacLennan DH. Isolation of a high affinity calcium-binding protein from sarcoplasmic reticulum. *J. Biol. Chem.* 249:974-9 (1974).
- Mesaeli N, et al. Calreticulin is essential for cardiac development. *J. Cell Biol.* 144:857-68 (1999).
- Michalak M, et al. Calreticulin, a multi-process calcium-buffering chaperone of the endoplasmic reticulum. *Biochem. J.* 417:651-66 (2009).
- Bedard K, Szabo E, Michalak M, Opas M. Cellular functions of endoplasmic reticulum chaperones calreticulin, calnexin, and ERp57. *Int. Rev. Cytol.* 245:91-121 (2005).
- Obeid M, et al. Calreticulin exposure dictates the immunogenicity of cancer cell death. *Nat. Med.* 13:54-61 (2007).
- Tesniere A, et al. Immunogenic cancer cell death: a key-lock paradigm. *Curr. Opin. Immunol.* 20:504-11 (2008).
- Gardai SJ, et al. Cell-surface calreticulin initiates clearance of viable or apoptotic cells through trans-activation of LRP on the phagocyte. *Cell* 123:321-34 (2005).
- Fucikova J, et al. Relevance of the chaperone-like protein calreticulin for the biological behavior and clinical outcome of cancer. *Immunol. Lett.* 193:25-34 (2018).
- Birgisdottir AB, Lamark T, Johansen T. The LIR motif - crucial for selective autophagy. *J. Cell Sci.* 126:3237-47 (2013).
- Yang Y, et al. The ER-localized Ca²⁺-binding protein calreticulin couples ER stress to autophagy by associating with microtubule-associated protein 1A/1B light chain 3. *J. Biol. Chem.* 294:772-782 (2019).
- Slobodkin MR, Elazar Z. The Atg8 family: multifunctional ubiquitin-like key regulators of autophagy. *Essays Biochem.* 55:51-64 (2013).

Applications	Host	Isotype	Molecular Wt.	Species Cross-Reactivity
WB, IF/ICC, IHC	Mouse	IgG1	48kDa	Hu, Rt, Ms



Western blot analysis of lysates of different cell lines using mouse mAb to calreticulin, MCA-6C6, dilution 1:2,000: [1] protein standard, [2] NIH-3T3, [3] HEK293, [4] HeLa, [5] SH-SY5Y, [6] C6, [7] COS-1 cells. A strong single band at about 50kDa corresponds to the calreticulin protein.



Immunofluorescent analysis of SH-SY5Y cells stained with mouse mAb to calreticulin, MCA-6C6, dilution 1:500 in green and costained with chicken pAb to lamin A/C, CPCA-LaminAC, dilution 1:2,000 in red. The blue is DAPI staining of nuclear DNA. The MCA-6C6 antibody reveals granular staining of cytoplasm, while the lamin A/C antibody stains the nuclear lamina and membrane.

Background:

Calreticulin was first identified as a calcium binding protein found in rabbit skeletal muscle. It appears to be a multifunctional protein found predominantly in endoplasmic reticulum (ER) and is ubiquitously expressed in a wide range of species (1). The protein has a conserved globular ~170 amino acid lectin-like N-terminal domain, a central proline rich region and a C-terminal low affinity but high capacity acidic Calcium binding region. There is a four-amino acid ER retention sequence (KDEL) at the extreme C-terminus accounting for ER localization. Gene knock out experiments show that the absence of the calreticulin gene is embryonically lethal (2). One of the major functions of calreticulin appears to be buffering Calcium levels in the ER which also affects intracellular calcium signaling (3). Calreticulin also functions as an autophagy receptor and as a molecular chaperone for glycoproteins (4). Calreticulin may also be found on the cell surface where it may mediate phagocytosis of apoptotic or dying tumor cells by triggering immune responses and allowing the immune system to remove tumor cells. This suggests that targeting cell surface calreticulin may be a novel anti-tumor therapy (5-8). The protein includes a functional "LC3 interacting region" or LIR motif (9). Proteins with LIR motifs bind ATG8 family members which in mammals are the LC3 and GABARAP proteins and calreticulin has been shown to bind binds LC3 (10). This binding is one of the essential steps in the process of autophagy, so LIR motif proteins generally function to target other proteins which bind to them for degradation.

The MCA-6C6 antibody was raised against the peptide VESGSLEDDWDFLPPKKI, corresponding to amino acids 191-208 of the human calreticulin protein. This peptide includes the LIR motif of the molecule, centered on the peptide WxxL, so this antibody may block binding of other molecules to this site (9,11). The antibody works well for western blotting and for IF, ICC and IHC (for IHC see data under "Additional Info" tab).

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.